

CORRECTION

Open Access



Correction to: *Helicobacter*-induced gastric inflammation alters the properties of gastric tissue stem/progenitor cells

Wataru Shibata^{1,2}, Soichiro Sue¹, Sachiko Tsumura^{1,3}, Yasuaki Ishii¹, Takeshi Sato¹, Eri Kameta¹, Makoto Sugimori¹, Hiroaki Yamada¹, Hiroaki Kaneko¹, Tomohiko Sasaki¹, Tomohiro Ishii¹, Toshihide Tamura¹, Masaaki Kondo¹ and Shin Maeda^{1*}

Correction

Unfortunately, the original article [1] contained an error incorporated during production. A duplicated version of Table 1 was published in place of Table 2. Table 2 has been corrected in the original article and is also included correctly below.

Author details

¹Department of Gastroenterology, Yokohama City University Graduate School of Medicine, Yokohama, Japan. ²Division of Translational Research, Advanced Medical Research Center, Yokohama City University, Yokohama, Japan. ³School of Medicine, Yokohama City University, Yokohama, Japan.

Published online: 08 January 2018

Reference

1. Shibata W, et al. *Helicobacter*-induced gastric inflammation alters the properties of gastric tissue stem/progenitor cells. *BMC Gastroenterol.* 2017; 17:145. <https://doi.org/10.1186/s12876-017-0706-6>.

Table 2 List of representative genes differentially expressed between organoids with and without *H. felis* infection

Gene Name	Log2 ratio
Intestine related genes	
<i>Isx</i>	2.75
<i>Muc4</i>	2.10
<i>Muc16</i>	1.70
<i>Vil1</i>	1.10
Gastric genes	
<i>Pepsinogen C</i>	- 3.17
<i>Gif</i>	- 2.47
Others	
<i>Gastrin</i>	3.30
<i>Cd34</i>	2.77
<i>Il-1a</i>	2.51
<i>Cd274 (Pd-I1)</i>	1.31

* Correspondence: smaeda@med.yokohama-cu.ac.jp

¹Department of Gastroenterology, Yokohama City University Graduate School of Medicine, Yokohama, Japan

